


How Accurately Do We Know the Formation of Solar ${}^8\text{B}$?

Moshe Gai ^{a *}

^aLaboratory for Nuclear Science at Avery Point, University of Connecticut, 1084 Shennecossett Rd., Groton, CT 06340. e-mail: moshe.gai@uconn.edu, URL: <http://www.phys.uconn.edu>

A detailed examination of current data on S_{17} (as opposed to an examination of $S_{17}(0)$ only) excludes quoting $S_{17}(0)$ with sufficiently small uncertainty. In contrast to suggestions that $S_{17}(0)$ is now known with the accuracy of $\pm 3\%$, the exact value of $S_{17}(0)$ is dependent on the choice of the data and the choice of theory used for extrapolation. In addition recent high precision results (including the Seattle data) on S_{17} which are in good agreement, still differ on the measured slopes, as do theoretical models that predict different d-wave contribution, precluding an accurate extrapolation to zero energy of the consistent data. Using a common extrapolation of only the consistent high precision data, suggests a value of $S_{17}(0) = 21.2 \pm 0.5$ eV-b, but a value equal to or smaller than 19.0 eV-b can not be excluded due to the uncertainty in the extrapolation, leading to an additional error of $^{+0.0}_{-3.0}$ eV-b. This (unacceptable) situation must be cleared by future experiments. .



*Work Supported by USDOE Grant No. DE-FG02-94ER40870.